

Vaccination Schedule

2024.11

☆ For routine vaccinations, please check the expiration date of the pre-vaccination questionnaire.

		Name of Vaccine (preventable Disease)	Type of vaccine	Age to begin	Recommended period and interval by our clinic
Before 1year of age	Routine	DPT-IPV-Hib (Diphtheria, Pertussis, Tetanus, Polio, Hib infection)	Inactivated	After 2months	1st dose : 2months of age 3doses in 4-8weeks interval booster : 1year or older (during 6-18months after the 3rd dose)
	Routine	PCV (pneumococcal infection)	Inactivated	After 2months	1st dose : 2months of age 3doses in 4weeks or more interval booster : 12-15 months of age (60days or more after the 3rd dose)
	Routine	HB (HepatitisB)	Inactivated	After 2months	1st dose : 2month of age 2nd dose : 4 or more weeks after the 1st dose 3rd dose : 20 or more weeks after the 1st dose ※mother-to-child transmission program (voluntary vaccine) 1st dose :0month 2nd dose :1month 3rd dose :6months (of age)
	Routine	Rota (Rotavirus gastroenteritis)	Live (oral)	After 6weeks	1st dose : 2months, just by 15weeks of age 2nd dose : 4 or more weeks after the 1st dose , just by 24weeks of age ※Another type of Rotavirus vaccine for 3-times vaccinations is available. It is not possible to receive different type of vaccine alternately.
	Routine	BCG (Tuberculosis)	Live (percutaneous)	After 3months	1 dose at 5~8months of age , just by 1year of age
After 1year of age	Routine	MR (Measles, Rubella)	Live	After 1year	Stage1 : 1 dose at 1year of age Stage2 : 1 dose in the year before admission into elementary school
	Routine	Chickenpox	Live	After 1year	1st dose : 1 year of age 2nd dose : during 6-12 months after the 1st dose , just by 3years of age
	Voluntary	Mumps	Live	After 1year	1st dose : 1 year of age 2nd dose : 5-6 years of age
	Routine	Japanese Encephalitis	Inactivated	After 3years	Stage1 : 1st dose:3years of age 2nd dose:1-4weeks after the 1st dose booster: 1year after the 2nd dose Stage2 : 1 dose at 9-12years of age ※Children at high risk of contracting JE can receive vaccination starting at 6 months of age.
	Voluntary	DPT (Diphtheria, Pertussis, Tetanus)	Inactivated	After 5years	5-6 years of age ※It is also acceptable to administer the DPT vaccine instead of the DT vaccine.
	Routine	DT (Diphtheria, Tetanus)	Inactivated	After 11years	1 dose at 11-12 years of age
	Routine	HPV 9-valent (Cervical cancer)	Inactivated	6th grade of elementary school(girls)	1st dose : 1st grade of junior high school 2nd dose : 6~12months after the 1st dose ※Age at the time of the 1st dose is 15 years or older, additional vaccinations are administered 2 months and 6 months after the 1st dose (total of 3 doses)
	Voluntary	Influenza (Seasonal influenza)	Inactivated	After 6months	2 doses in 2-4 weeks interval (yearly Oct.-Dec.) ※1 dose for the 13years of age or older
			Live (nasal)	After 2years	1 dose at 2-18 years of age (yearly Oct.-Dec.)